

中国原哈格鸣蠡科化石新发现 (直翅目, 哈格鸣蠡科, 阿博鸣蠡亚科)

李连梅 任 东* 孟祥明

首都师范大学生命科学学院 北京 100037

摘 要 描述原哈格鸣蠡科化石 3 新属, 3 新种: *Flexaboilus retinervius* gen. et sp. nov., *Angustaboilus fangianus* gen. et sp. nov., *Novaboilus multifurcatus* gen. et sp. nov.。所有化石标本均采自于内蒙古宁城道虎沟中侏罗世九龙山组地层, 现保存于首都师范大学生命科学学院。

关键词 直翅目, 原哈格鸣蠡科, 阿博鸣蠡亚科, 新属, 新种, 中国。

中图分类号 Q915.819.7

阿博鸣蠡亚科 Aboilinae 是目前原哈格鸣蠡科已发表的 6 亚科之一。Martynov 于 1925 年建立了阿博鸣蠡科 Aboilidae, 后来 Gorochoy (1986) 将其调整为阿博鸣蠡亚科 Aboilinae。Aboilinae 很可能是 Protaboilinae 的后裔, 它是 Prophalangopsidae 最为丰富和多样性的亚科 (Gorochoy, 2003)。

Aboilinae 目前已发表 13 属, 主要分布年代为中侏罗世和晚侏罗世, 主要分布地区为俄罗斯西伯利亚、中国、哈萨克斯坦及蒙古国 (Gorochoy, 2003)。

最近作者自内蒙古宁城道虎沟九龙山组地层采得大量昆虫化石标本, 其中包括了本文描述的 3 件阿博鸣蠡亚科化石。其化石归属于中侏罗世九龙山组 (任东, 2002)。本文描述的标本保存于首都师范大学昆虫演化与变迁重点实验室。

本文所有线条图均借助于 Leica 显微镜附带绘图臂完成。采用的术语参考 Gorochoy 所采用的系统。

原哈格鸣蠡科 Prophalangopsidae Kirby, 1906

阿博鸣蠡亚科 Aboilinae Martynov, 1925

Aboilidae Martynov, 1925. *Bull. Acad. Sci. Russie*, 569-598.

Aboilinae Gorochoy, 1986. *Trudy Zool. Inst. Akad. Nauk SSSR*, 143: 65-100.

Aboilinae Gorochoy, 1988. *Paleontol. Zhurnal*, (2): 54-66.

Aboilinae Gorochoy, 1995. Pt. 1. *Trudy Zool. Inst. Ross. Akad.*, 260: 126-137.

弯曲阿博鸣蠡属, 新属 Flexaboilus gen. nov.

模式种: *Flexaboilus retinervius* sp. nov.

词源: flex- (源自拉丁词 flex-, 弯曲) + aboilus (Aboilinae 亚科 aboilus 属), 阳性。

鉴别特征 雄虫前翅 Fc 脉短, 呈弧形, 前缘域

狭长。Sc 脉略呈 S 形, 支脉多于 15 枝。R 脉在翅长 1/3 前分支, 支脉 5 枝。Rs 脉支脉 9 枝。M-Cu 域基干部与 Sc-R 域基干部近等宽。MP+CuA₁ 脉与 R 脉同时分支, 支脉 6 枝。2CuA₂ 较倾斜, 不与 1CuA₂ 平行。CuP 弯曲不强烈。

雌虫: 未知。

比较与讨论 依据前翅 Fc 脉横切 Sc 脉支脉, MA 脉不与 Rs 脉汇合, MA 脉在 R-M 域最宽处分支, 支脉 2 枝, MP 脉与 CuA₁ 脉融合, 合并一段后分支等特征将新属归于原哈格鸣蠡科。又由 Fc 脉较发达, MP+CuA₁ 支脉未到达翅顶角将其归入阿博鸣蠡亚科。

本亚科目目前已发表 13 属 (Gorochoy, 2003), 其中 *Aboilus* Martynov, 1925 分布于哈萨克斯坦和俄罗斯西伯利亚地区晚侏罗世; *Sinoprophanlangopsis* Hong, 1982 分布于中国中侏罗世; *Bacharaboilus* Gorochoy, 1988 分布于蒙古国中侏罗世; *Apsataboilus* Gorochoy, 1990 分布于俄罗斯西伯利亚晚侏罗世和早白垩世; *Prophalangopsides* Sharov, 1968 分布于俄罗斯西伯利亚早白垩世; *Tettaboilus* Gorochoy, 1988 分布于俄罗斯西伯利亚早白垩世; *Utanaboilus* Gorochoy, 1990 分布于俄罗斯西伯利亚早白垩世也可能是晚白垩世; *Brunnus* Hong, 1983 分布于中国中侏罗世; *Pseudohagla* Sharov, 1962 分布于俄罗斯西伯利亚中侏罗世到晚侏罗世; *Rynophlebia* Deichmüller, 1886 分布于德国晚侏罗世; *Nipponohagla* Fujiana, 1978 分布于日本早白垩世; *Baissaboilus* Gorochoy, 1996 分布于俄罗斯西伯利亚早白垩世; *Karatailus* Gorochoy, 1996 分布于哈萨克斯坦晚侏罗世。

国家自然科学基金 (30025006, 30370184, 30430100)、北京市自然科学基金 (5032003)、北京市教育委员会科技发展计划重点项目 (KZ200410028013) 和北京市属市管高校人才强教计划共同资助。

* 通讯作者, E-mail: rendong@mail.cnu.edu.cn

收稿日期: 2006-10-10, 修订日期: 2006-10-26.

新属与已知属差异较显著，前翅 Fc 脉短，前缘域狭长；R₁ 脉，Rs 脉及 MP+ CuA₁ 脉分支均较早，分支较多；2CuA₂ 较倾斜，不与 1CuA₂ 平行。

新属与 *Abolilus* Martynov, 1925 较为相近，但新属前缘域狭长，2CuA₂ 较倾斜，不与 1CuA₂ 平行，R 脉与 MP+ CuA₁ 脉同时分支，CuP 弯曲不强烈。

网脉弯曲阿博鸣蠹，新种 *Flexaboilus retinervius* sp. nov. (图 1, 6)

正模：一块保存较完整的雄虫前翅标本，编号：CNU O-NN 2006107。

词源：“*retinervius*”源自拉丁词，网状脉的。

产地及层位：内蒙古宁城道虎沟，九龙山组（中侏罗世）。

描述 雄虫前翅翅长 71 mm，翅宽 33 mm（保存部分）。翅面具深色带斑（见图 6）。翅基部较宽，近端部逐渐收缩。Fc 脉短，呈弧形，在翅长 1/4 处

达翅前缘。前缘域狭长，亚前缘域宽阔。Sc 脉略呈 S 形，在翅长 4/5 处融于前缘，支脉 16 枝，近平行排列，其间有间插脉。R 脉自翅基部发出，在翅长 1/3 稍前分支，R₁ 脉在超过翅长 1/3 后再分支，支脉 5 枝，近平行排列，其间规则横脉发育。Rs 脉在 R₁ 脉分支后稍后分支，支脉 9 枝，梳状排列，其间不规则横脉发育。Sc-R 域基干部较窄，其间规则横脉发育。R₁-Rs 域基干部较窄，其间规则横脉发育。M 脉自翅基部发出，MA 脉在 R 脉分支前叉状分支，2 支脉弓形，近平行排列，其间 2 组横脉发育。R-M 域基干部较宽，其间 2 组横脉发育。M-Cu 域基干部与 Sc-R 域基干部近等宽。MP 脉自 MA 脉分支前分出，与 CuA 脉融合，合并一段后分出。MP+ CuA₁ 脉与 R 脉同时分支，末端 6 枝，其间不规则横脉发育。2CuA₂ 较倾斜，不与 1CuA₂ 平行。CuP 近端较倾斜，远端呈弓形。




图 1 网脉弯曲阿博鸣蠹，新种 *Flexaboilus retinervius* sp. nov.，正模，雄性（holotype, male），CNU O-NN 2006107，前翅翅脉（draw line of the forewing）

狭阿博鸣蠹，新属 *Angustaboilus* gen. nov.

模式种：房氏狭阿博鸣蠹 *Angustaboilus fangianus* sp. nov.

词源：angust-（源自拉丁词 *angusti-*，狭）+ *aboilus*（*Abolilinae* 亚科 *aboilus* 属），阳性。

鉴别特征 雌虫前翅 Fc 脉短，呈弧形。前缘域略呈三角形，其间纵脉放射状排列。Sc 脉直，约在翅长 3/4 处达前缘，支脉多于 10 枝。R 脉约在翅长 1/3 处分支，支脉 7 枝。Rs 脉分支较晚，在 R₁ 脉第 2 支脉稍后分支，支脉 7 枝。M-Cu 域基干部与 Sc-R 域基干部近等宽。MP+ CuA₁ 脉与 R 脉同时分支，支脉 3 枝。

雄虫：未知。

比较与讨论 新属与 *Sinoprophanlangopsis* Hong,

1982 较为相似，但新属 Fc 脉成弧形，R₁ 脉与 Rs 脉分支较多，Rs 脉分支较晚等特征与其区别明显。该属与 *Utanaboilus* Gorochov, 1990 较为相近，但新属 R₁ 脉分支较多，MP+ CuA₁ 脉分支较少，Rs 脉分支较晚等特征与其区别明显。

房氏狭阿博鸣蠹，新种 *Angustaboilus fangianus* sp. nov. (图 2~4, 8~11)

正模：一块前后翅均保存的雌虫标本，编号：CNU O-NN 2006108。

词源：种名以房良先生的姓氏命名，感谢他赠与的化石。

产地及层位：内蒙古宁城道虎沟，九龙山组（中侏罗世）。




图 2~ 4 美丽狭阿博鸣螽, 新种 *Angustaboilus fangianus* sp. nov.

正模, 雌性 (holotype, male), CNU-ONN 2006108 2 左前翅 (left forewing) 3 右前翅 (right forewing) 4 后翅 (hind wing)

描述 雌虫前翅翅长 57 mm, 左翅翅宽 15 mm, 右翅翅宽 16 mm。翅面狭长, 呈深褐色 (图 8~ 11)。Fc 脉短, 呈弧形, 在翅长 1/2 稍前达翅前缘。前缘域狭长, 略呈三角形。亚前缘域宽阔。Sc 脉直, 约在翅长 3/4 处融于前缘, 左翅支脉 11 枝, 右翅支脉 12 枝, 近平行排列, 其间不规则横脉发育。R 脉自翅基部发出, 约在翅长 1/3 处开始分支, R₁ 脉在越

过翅长 1/3 处再分支, 左翅支脉 8 枝, 右翅支脉 7 枝, 近平行排列, 其间规则横脉发育。R_s 脉在 R₁ 脉第 2 支脉稍后分支, 支脉 7 枝, 梳状排列, 其间不规则横脉发育。Sc-R 域基干部较窄, 其间规则横脉发育。R₁-R_s 域基干部较宽, 其间有规则横脉发育。M 脉自 R 脉基部发出, MA 脉在 R 脉分支前叉状分支, 2 支脉弓形, 近平行排列, 其间有 2 组横脉

发育。R-M 域基干部较窄，其间有 2 组横脉发育。M-Cu 域基干部与 Sc-R 域基干部近等宽。MP 脉自 MA 脉分支前分出，与 CuA 脉融合，合并一段后分出。MP+CuA₁ 脉与 R 脉同时分支，末端 3 枝，近平行排列，其间规则横脉发育。1CuA₂ 脉与 2CuA₂ 脉，CuP 脉平行。1CuA₂-2CuA₂ 域与 2CuA₂-CuP 域较窄，其间规则横脉发育。A 脉 4 枝，其间规则横脉发育。

后翅翅长 45 mm，翅宽 15.5 mm（保存部分）。翅面呈深褐色（图 11）。翅基部强烈收缩。Sc 脉直，在翅长 1/2 稍后达翅前缘，支脉 12 枝，近平行排列。R 脉在翅长 1/5 处分支，R₁ 脉在翅长 1/3 处再分支，支脉 7 枝，近平行排列，其间规则横脉发育。Rs 脉在 R₁ 脉分支前分支，支脉 8 枝，近梳状排列，其间不规则横脉发育。MA 脉在 R 脉分支前叉状分支，两支脉平行排列，其间 2 组横脉发育。MP 自 M 脉基部发出，支脉 2 枝，其间规则横脉发育。

新阿博鸣蠹，新属 *Novaboilus* gen. nov.

模式种：多叉新阿博鸣蠹 *Novaboilus multifurcatus* sp. nov.

词源：nov（源自拉丁词 *novi*，新）+ aboilus（Aboilinae 亚科 *aboilus* 属），阳性。

鉴别特征 雄虫前翅 Fc 脉较短，呈弧形。前缘域较宽。Sc 脉直，在翅长 2/3 处达前缘，支脉少于 10 枝，其间支脉再分支。R 脉约在翅长 1/3 处分支，R₁ 脉支脉 4 枝。Rs 脉在 R₁ 脉分支前分支，支脉 8 枝。M-Cu 域基干部与 Sc-R 域基干部近等宽，MP+CuA₁ 脉分支略早于 R 脉，支脉 5 枝。

雌虫：未知。

比较与讨论 新属与 *Aboilus* Martynov, 1925 较为相近，但新属 Sc 脉分支较少，Rs 脉分支早于 R₁

脉，MP+CuA₁ 脉末端带 5 支等特征与其区别明显。新属与 *Brumaus* Hong, 1983 较为相近，但新属 Fc 脉短；前缘域较宽；Rs 脉，MP+CuA₁ 脉分支较多；R₁ 脉，Rs 脉分支点距离 R 脉分支点较近等特征与其区别明显。

多叉新阿博鸣蠹，新种 *Novaboilus multifurcatus* sp. nov.（图 5, 7）

正模，一件保存较好的雄虫前翅标本，编号：CNU G NN 2006090。

词源：“*multifurcatus*”源自拉丁词，多叉。

产地及层位：内蒙古宁城，九龙山组（中侏罗世）。

描述 雄虫前翅翅长 50 mm，翅宽 18 mm（保存部分）。翅面具深色带斑（图 7）。Fc 脉直，在翅长 1/2 稍前融于前缘，前缘域较宽。Sc 脉直，约在翅长 2/3 处达翅前缘，支脉 9 枝，其支脉再次分支，其间有间插脉发育。R 脉自翅基部发出，在翅长 1/3 处分支，R₁ 脉在翅长 1/2 处再次分支，支脉 4 枝，其间规则横脉发育。Rs 脉在 R₁ 脉分支前分支，支脉 8 枝，近梳状排列，其间不规则横脉发育。Sc-R 域基干部较宽，其间规则横脉发育。R₁-Rs 域基干部较宽，其间规则横脉发育。M 脉自翅基部发出，MA 脉在 R 脉分支前叉状分支，2 支脉弓形，近平行排列，其间规则横脉发育。R-M 域基干部较宽，其间规则横脉发育。M-Cu 域基干部与 Sc-R 域基干部近等宽。MP 脉自 MA 脉分支前分出，与 CuA 脉融合，合并一段后分出。MP+CuA₁ 分支稍早于 R 脉，末端 5 枝，其间规则横脉发育。2CuA₂ 较倾斜，不与 1CuA₂ 平行。MP+CuA₁-1CuA₁ 域及 1CuA₁-2CuA₂ 域较窄，其间规则横脉发育。




图 5 多叉新阿博鸣蠹，新种 *Novaboilus multifurcatus* sp. nov.，正模，雄性（holotype, male），CNU G NN 2006090，前翅翅脉（draw line of the forewing）



图 6 网脉弯曲阿博鸣螽，新种 *Flexaboilus retinerviis* sp. nov.

图 7 多叉新阿博鸣螽，新种 *Novaboilus multifurcatus* sp. nov.



图 8~ 11 美丽狭阿博鸣蠹，新种 *Angustaboilus fengianus* sp. nov.

8. 整体 (body with wings) 9. 左前翅 (left forewing) 10. 右前翅 (right forewing) 11. 后翅 (hind wings)

致谢 在研究工作中河北大学生命科学学院石福明教授提供了大量的资料，论文的完成得到了王莹，

王田田，刘明，孙建海，谭京晶，刘玉双，黄建东等同学的帮助，在此表示感谢。

REFERENCES (参考文献)

- Gorochov, A. V. 1986. Triassic insects of the Superfamily Hagloidea (Orthoptera). Trudy Zool. Inst. Akad. Nauk SSSR 143 Zool. Inst. Akad. Nauk SSSR, Leningrad. 65-100. (in Russian).
- Gorochov, A. V. 1988. The Lower And Middle Jurassic Superfamily Hagloidea (Orthoptera). *Palcont. Jav.*, 2: 50-61.
- Gorochov, A. V. 1990. Orthoptera Gryllida. In: Rasnitsyn, A. P. (ed.), Late Mesozoic insects of Eastern Transbaikalia. Trudy Palcont. Inst. Akad. Nauk SSSR 239 Nauka, Moscow. 210-214. (in Russian).
- Gorochov, A. V. 1995. System and Evolution of the Suborder Ensifera (Orthoptera). Pt. 1. Trudy Zool. Inst. Ross. Akad. Nauk 260. Zool. Inst. Ross. Akad. Nauk, St. - Petersburg. 126-137. (in Russian).
- Gorochov, A. V. 1996. New Mesozoic insects of the Superfamily Hagloidea (Orthoptera). *Palcont. Zhurnal* (3): 73-82 (in Russian).
- Gorochov, A. V. 2003. New data on taxonomy and evolution of fossil and Recent Prophalangopsidae (Orthoptera: Hagloidea). *Acta Zoologica Graecovinsia*, 46 (suppl. - Fossil Insects): 117-127.
- Hong, Y-C 1982a. Fossil Haglidae (Orthoptera) in China. *Scientia Sinica* (series B), 25 (10): 1118-1129. [洪友崇, 1982a. 中国直翅目哈格鸣螽科化石. 中国科学 B 辑, 25 (10): 1118-1129]
- Hong, Y-C 1982b. Mesozoic Fossil Insects of Jiuquan Basin in Gansu Province. Geological Publishing House, Beijing. 7-80. [洪友崇, 1982b. 酒泉盆地昆虫化石. 北京: 地质出版社. 71-80]
- Hong, Y-C 1983. Middle Jurassic Fossil Insects in North China. Geological Publishing House, Beijing. 42-48. [洪友崇, 1983. 北方中侏罗世昆虫化石. 北京: 地质出版社. 42-48]
- Hong, Y-C 1984. II. Mesozoic Volume. In: Palaeontological Atlas of North China. Geological Publishing House, Beijing. 148-152. [洪友崇, 1984. (二) 中生代分册. 华北地区古生物图册. 北京: 地质出版社. 148-152]
- Hong, Y-C 1986. New fossil insects of Haifanggou Formation, Liaoning Province. *J. Changchun Coll. Geol.*, 4: 10-16. [洪友崇, 1986. 辽西海房沟组新的昆虫化石. 长春地质学院学报, 4: 10-16]
- Hong, Y-C 1990. The stratigraphy and palaeontology of Laiyang Basin, Shandong Province. Geological Publishing House, Beijing. 69-71. [洪友崇, 1990. 山东莱阳盆地地层古生物. 北京: 地质出版社. 69-71]
- Hong, Y-C 1992. Middle and Late Jurassic Fossil Insects. In: Palaeontological Atlas of Jilin Province. Jilin Science and Technology Publishing House, Changchun. 420-421. [洪友崇, 1992. 中晚侏罗世昆虫. 吉林省古生物图册. 长春: 吉林科技出版社. 420-421]
- Lin, Q-B 1965. Two insects from the Lower Part of Jurassic, Inner Mongolia. *Acta Palaeont. Sin.*, 13 (2): 363-368. [林启彬, 1965. 内蒙下部侏罗世的两种昆虫化石. 古生物学报, 13 (2): 363-368]
- Martynov, A. V. 1925. To the knowledge of fossil insects from Jurassic beds in Turkestan. *Bull. Acad. Sci. Russie*, 569-598.
- Ren, D 1995. Fossil Insects Part. In: Fauna and Stratigraphy of Jurassic Cretaceous in Beijing and the Adjacent Areas. Earthquake Publishing House, Beijing. 61-65. [任东, 1995. 昆虫化石部分. 北京与邻区侏罗-白垩纪动物群及其地层. 北京: 地震出版社. 61-65]
- Ren, D 2002. Stratigraphic division of the Jurassic in the Daohugou area, Ningcheng, Inner Mongolia. *Geological Bulletin of China. Sinica*, 21 (8-9): 584-591. [任东, 2002. 内蒙古宁城道虎沟地区侏罗纪地层划分及时代探讨. 地质通报, 21 (8-9): 584-591]
- Sharov, A. G. 1968. Phylogeny of the Orthopteroidea. Trudy Palcont. Inst. Akad. Nauk SSSR 118. Nauka, Moscow. 218 p. (in Russian)

NEW FOSSIL PROPHALANGOPSIDS FROM CHINA (ORTHOPTERA, PROPHALANGOPSIDAE, ABOILINAE)

LI Lian-Mei, REN Dong, MENG Xiang-Ming

College of Life Science, Capital Normal University, Beijing 100037, China

Abstract In this paper three new genera and three new species were described: *Flexaboilus retinervius* gen. et sp. nov., *Angustaboilus fangianus* gen. et sp. nov. and *Novaboilus multifurcatus* gen. et sp. nov. All type specimens were collected from the Middle Jurassic Jiulongshan Formation at Daohugou Village, Ningcheng, Inner Mongolia, China and deposited in Capital Normal University.

Flexaboilus gen. nov.Holotype: *Flexaboilus retinervius* sp. nov.Etymology. Flex- (from Latin *flex*, means bend) + aboilus (name of genus *aboilus*), masculine.

Diagnosis. Male, Fc short, arched. Precostal field narrow. Sc S-shaped, with more than 15 branches. R forked before 1/3 of the wing length, with 5 branches. Rs with 9 branches. Width of the basal part of M-Cu field equal to the basal part of Sc-R field. MP + CuA₁ and R forked at the same level, with 6 branches. 2CuA₂ quite incline, and not parallel to 1CuA₂. CuP slightly curve.

Female: unknown.

Discussion. The new genus was assigned to the family Prophalangopsidae by elytrum with branches of Sc

intersected by developed Fc; MA not fused with Rs and branched off from most convex part of R-M field, with 2 branches; MP coalesced with CuA₁ basally and then diverged again.

The new genus was assigned to the subfamily Aboilinae by well developed Fc, MP + CuA₁ ending considerably before reaching vertex of elytrum.

At present, the subfamily consists of 13 previously described extinct genera (Gorochov, 2003). *Aboilus* Martynov, 1925 from Upper Jurassic, Kazakhstan and Siberia; *Sanoprophanlangopsis* Hong, 1982 from middle Jurassic, China; *Budharaboilus* Gorochov, 1988 from Middle Jurassic, Mongolia; *Apsataboilus* Gorochov, 1990 from Upper Jurassic or Lower Cretaceous, Siberia; *Prophalangopsides* Sharov, 1968 from Lower Cretaceous, Siberia; *Tettaboilus* Gorochov, 1988 from Lower Cretaceous, Siberia; *Utanaboilus* Gorochov, 1990 from Lower and Possibly Upper Cretaceous, Siberia; *Brumeus* Hong, 1983 from middle Jurassic, China; *Pseudohagla* Sharov, 1962 from Middle-Upper Jurassic of Siberia; *Bynophlebia* Deichmuller, 1886 from Upper Jurassic of Germany; *Nipponohagla* Fujiama, 1978 from Lower Cretaceous of Japan; *Baissaboilus* Gorochov, 1996 from

Lower Cretaceous of Siberia; *Karatailus* Gorochoy, 1996 Upper Jurassic, Kazakhstan. The new genus different from them by Fc short; precostal field narrow; R_1 , R_s and $MP+CuA_1$ beginning to branch very early, with numerous branches; $2CuA_2$ slightly curve, and not parallel to $1CuA_2$.

The new genus is similar to *Aboilus* Martynov, 1925, but were different from it in following characters: Precostal field narrow; $2CuA_2$ quite incline and not parallel to $1CuA_2$; R and $MP+CuA_1$ forked at the same level and CuP slightly curve.

Flexaboilus retinervius **sp. nov.** (Figs 1, 6)

Etymology. “*retinervius*” from Latin, means meshwork.

Locality and horizon: Daohugou Village, Ningcheng, Inner Mongolia. Middle Jurassic, Jiulongshan Formation.

Description. Forewing length 71 mm, width 33 mm (as preserved). Fc short, arched. Sc S-shaped, with 16 parallel branches. R forked before $1/3$ of the wing length, with 5 parallel branches. R_s forked later than R_1 , with 9 parallel branches. $MP+CuA_1$ and R forked at the same level, with 6 branches. $2CuA_2$ quite incline, and not parallel to $1CuA_2$. CuP slightly curve.

Angustaboilus **gen. nov.**

Holotype: *Angustaboilus fangianus* sp. nov.

Etymology. angust- (from Latin *angusti-*, means narrow) + *aboilus* (name of genus *aboilus*), masculine.

Diagnosis. Female, Fc short, arched. Precostal field triangle, veinlets radial. Sc straight, extending about $3/4$ of the wing length, with more than 10 branches. R forked at $1/3$ of the wing length, with 7 branches. R_s forked after second branch of R_1 , with 7 branches. Width of the basal part of $M+Cu$ field equal to the basal part of $Sc+R$ field. $MP+CuA_1$ and R forked at the same level, with 3 branches.

Male: unknown.

Discussion. This new genus is similar to *Sinoprophanlangopsis* Hong, 1982, but were different from the latter by following characters: Fc arched, R_1 and R_s with several branches and R_s forked very late. The new genus is also similar to *Utanoaboilus* Gorochoy, 1990, however, it could be distinguished from the latter by following characters: R_s with numerous branches, $MP+CuA_1$ with few branches and R_s branched very late.

Angustaboilus fangianus **sp. nov.** (Figs 2-4, 8-11)

Etymology. “*fangianus*” was name after Fang Liang for the fossil that he donated.

Locality and horizon. Daohugou Village,

Ningcheng, Inner Mongolia. Middle Jurassic, Jiulongshan Formation.

Description. Forewing length 57 mm, width 15 mm. Fc short, arched. Precostal field triangle, veinlets radial. Sc straight, extending at about $3/4$ of the wing length, left forewing with 11 branches, right forewing with 12 branches. R forked at $1/3$ of the wing length, left forewing with 8 branches, right forewing with 7 branches. R_s forked after second branch of R_1 , with 7 branches. $MP+CuA_1$ and R forked at the same level, with 3 branches. Hind wing length 45 mm, width 15.5 mm (preserved). Sc straight, with 12 branches. R forked at $1/5$ of the wing length, with 7 parallel branches. R_s forked earlier than R , with 8 parallel branches.

Novaboilus **gen. nov.**

Holotype: *Novaboilus multifurcatus* sp. nov.

Etymology. nov- (from Latin *novi-*, means new) + *aboilus* (name of genus *aboilus*), masculine.

Diagnosis. Male, Fc short, arched. Precostal field broad. Sc straight, extending at about $2/3$ of the wing length, with less than 10 branches, the branches forked again. R forked at $1/3$ of the wing length. R_1 with 4 branches. R_s forked earlier than R_1 , with 8 branches. Width of the basal part of $M+Cu$ field equal to the basal part of $Sc+R$. $MP+CuA_1$ forked earlier than R , with 5 branches.

Female: unknown.

Discussion. This new genus is similar to *Aboilus* Martynov, 1925, but were different from it in following characters: Sc with few branches, R_s forked earlier than R_1 and $MP+CuA_1$ with 5 branches. The new genus is also similar to *Brumeus* Hong, 1983. Whereas the new genus could be distinguished from the latter by Fc short, precostal field broad, R_s and $MP+CuA_1$ with numerous branches, R_1 and R_s beginning to branch very far from their bases.

Novaboilus multifurcatus **sp. nov.** (Figs 5, 7)

Etymology. “*multifuratus*” from Latin, means many forks.

Locality and horizon: Daohugou Village, Ningcheng, Inner Mongolia. Middle Jurassic, Jiulongshan Formation.

Description. Forewing length 57 mm, width 15 mm (as preserved). Fc short, arched. Sc straight, extending at about $2/3$ of the wing length, with 9 branches. R forked at $1/3$ of the wing length, with 4 branches. R_s forked earlier than R_1 , with 8 parallel branches. $MP+CuA_1$ forked earlier than R , with 5 branches.

Key words Orthoptera, Prophalangopsidae, Aboilinae, new genus, new species, China.